ISEMBYLD™
(apitegromab-mstn) injection 150mg/3mL

WHAT TO EXPECT WITH TREATMENT



Your guide to the infusion process

What is ISEMBYLD (apitegromab-mstn)?

ISEMBYLD is a prescription medicine used to treat spinal muscular atrophy (SMA) in adults and children 2 years of age and older who are currently receiving a survival motor neuron 2 (SMN2)-targeted treatment.

IMPORTANT SAFETY INFORMATION

Before taking ISEMBYLD, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of low bone density or bone fractures
- are pregnant or plan to become pregnant. If you are pregnant or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. It is not known if ISEMBYLD will harm your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with ISEMBYLD
- are breastfeeding or plan to breastfeed. It is not known if ISEMBYLD passes into breast milk. Talk to your healthcare provider about the best way to feed your baby while on treatment with ISEMBYLD

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

ISEMBYLD™ (apitegromab-mstn) injection 150mg/3mL

Getting started with ISEMBYLD

In choosing ISEMBYLD, you've made an important decision in your spinal muscular atrophy (SMA) journey. ISEMBYLD is the first and only FDA-approved muscle-targeted treatment for SMA. **ISEMBYLD is approved for adults and children 2 years of age and older who are currently receiving a survival motor neuron 2 (SMN2)-targeted treatment.**

When used with an SMN2-targeted treatment, ISEMBYLD can help you gain more motor function than if you were on an SMN2-targeted treatment alone.

This brochure is designed to help you understand the basics of ISEMBYLD infusion treatments and what you need to know before, during, and after each infusion.

INSIDE THIS BROCHURE

AN INTRODUCTION TO ISEMBYLD	3
PLANNING FOR ISEMBYLD	4
PREPARATION TIPS	5
INFUSION DAY: WHAT TO EXPECT	6
FAQS	7
ABOUT SCHOLAR ROCK SUPPORTS	8
IMPORTANT SAFETY INFORMATION	9

ScholarRock SUPPORTS™

If you have questions at any time, help is available through Scholar Rock Supports Monday – Friday 8 AM – 8 PM ET at 1-833-777-5444 or at ScholarRockSupports.com where you can find a dedicated **Rx Onboarding and Care Coordinator (ROCC)**.

IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider, including your pharmacist, when you get a new medicine.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

ISEMBYLD™ (apitegromab-mstn) injection 150mg/3mL

ISEMBYLD infusion overview

ISEMBYLD is a prescription medicine used to treat SMA in adults and children 2 years of age and older who are currently receiving a survival motor neuron 2 (SMN2)-targeted treatment. **ISEMBYLD is administered once every 4 weeks as an intravenous (IV) infusion. IV infusion is a standard procedure administered by a healthcare professional.**

INFUSION SCHEDULE	You'll receive an IV infusion of ISEMBYLD every 4 weeks
INFUSION DURATION	Your ISEMBYLD infusion should last approximately 1-2 hours
CUSTOMIZED TO YOU	Your ISEMBYLD dose will be based on your weight (10 mg/kg)
CONVENIENT LOCATIONS	ISEMBYLD is administered by a healthcare professional. Based on eligibility, infusions can be given at convenient locations, including: • Hospital • Home • Infusion center

IMPORTANT SAFETY INFORMATION

What are the possible side effects of ISEMBYLD?

- **Fractures.** Treatment with ISEMBYLD may increase the risk of bone fractures, including serious fractures. Bone fractures may happen with or without a fall or other injury. Your healthcare provider may consider stopping treatment with ISEMBYLD if you experience a bone fracture during treatment

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

ISEMBYLD™ (apitegromab-mstn) injection 150mg/3mL

Planning for your ISEMBYLD infusion

When you're ready for your first infusion, Scholar Rock Supports can help guide the way. You will have access to personalized support through a dedicated **Rx Onboarding and Care Coordinator (ROCC)** who will be there to support you at every step of your treatment journey. Your ROCC can also connect you to a **Regional Clinical Specialist (RCS)** to provide education and support on infusions, regardless of whether performed in a hospital, home, or infusion center.

Scheduling

Your ISEMBYLD treatment plan will consist of 1 infusion every 4 weeks. When scheduling, choose a time and day of the week that will work best for you on a regular basis, and save yourself some time by scheduling a few infusions in advance. You can work with your ROCC to get support when setting up your infusions.

Going to a Hospital or Infusion Center

It's important to factor in travel time if you're going to a hospital or infusion center. Consider saving the travel instructions on your phone or printing them out. Be sure to allow time to check in. Arriving early can make for a more comfortable infusion experience for everyone.

Getting Ready for a Home Infusion

Your ROCC can help connect you with your home infusion specialist or pharmacy to share what you need to know to prepare for a home infusion — including delivery instructions and guidance on how and where to store your treatment.

IMPORTANT SAFETY INFORMATION

The most common side effects of ISEMBYLD include:

- upper respiratory tract infections
- vomiting
- cough
- viral infections
- headache
- stomach flu (gastroenteritis)
- sore throat (pharyngitis)
- hypersensitivity

These are not all of the possible side effects of ISEMBYLD. Call your healthcare provider for medical advice about side effects.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

ISEMBYLD™ (apitegromab-mstn) injection 150mg/3mL

Getting ready for infusion day

Over time, the infusion process will become a regular routine. But these tips may help as you prepare for your first few infusions. You can speak with your **Rx Onboarding and Care Coordinator (ROCC)** before your first infusion for more information on how to prepare. If you have additional questions about the infusion process, your ROCC can connect you to a Regional Clinical Specialist (RCS).

Considerations for what to have for your infusion:

- A list of medications you are taking
- A list of the healthcare providers who manage your care
- A pillow and blanket
- Food or snacks
- A bottle of water to help stay hydrated (Note: it's important to increase water intake 24 hours before your infusion to improve access to veins)
- Chargers for any devices you want to use
- Something to read, play, or watch to pass the time—with headphones or earbuds, so you don't disturb others
- To stay comfortable, consider wearing loose-fitting clothes and dressing in layers

Ask if a friend or companion can be with you. **And if you have any other questions about your infusion, you can call your ROCC. They'll be happy to help.**

Mindy
Treated with ISEMBYLD

IMPORTANT SAFETY INFORMATION

Before taking ISEMBYLD, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of low bone density or bone fractures
- are pregnant or plan to become pregnant. If you are pregnant or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. It is not known if ISEMBYLD will harm your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with ISEMBYLD
- are breastfeeding or plan to breastfeed. It is not known if ISEMBYLD passes into breast milk. Talk to your healthcare provider about the best way to feed your baby while on treatment with ISEMBYLD

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

ISEMBYLD™ (apitegromab-mstn) injection 150mg/3mL

Infusion Day: What to expect, step by step

Before Infusion

Before starting, the nurse or other healthcare professional will go over the infusion process with you and answer any questions. They will also check and record your vital signs, such as pulse, temperature, and blood pressure, and may ask your weight to ensure you're getting the correct dosage.

During Infusion (approximately 1-2 hours)

An IV infusion is given through a needle into a vein in your arm or back of your hand. The needle is attached to a small tube connected to the infusion bag that contains the appropriate dose of ISEMBYLD based on your weight (10 mg/kg).

Once your infusion has begun, it's okay to stay occupied with books, tablets, or small toys, or to simply rest.

Your nurse or other healthcare professional will monitor you closely from start to finish to ensure that you're comfortable.
They will watch out for any side effects related to your infusion.

After Infusion

When your infusion is complete, your nurse or other healthcare professional will remove the needle, clean and bandage the area, and let you know when your visit is over.

You may be asked to schedule a time for your next infusion in 4 weeks if it has not already been scheduled.

If You Miss a Dose of ISEMBYLD

- ISEMBYLD should be given (administered) as soon as possible. Then, restart the dosing schedule 4 weeks after the missed dose is given, or
- The dose can be skipped and the next dose can be given as originally scheduled

IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider, including your pharmacist, when you get a new medicine.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

ISEMBYLD™ (apitegromab-mstn) injection 150mg/3mL

Frequently asked questions

What are the most common side effects?

The most common side effects (reported in at least 20% of participants treated with ISEMBYLD and more frequently than in participants on an SMN2-targeted treatment alone) were upper respiratory tract infections, vomiting, cough, viral infections, headache, stomach flu (gastroenteritis), sore throat (pharyngitis), and hypersensitivity. Any adverse events should be reported to your doctor.

How can I find an infusion center?

Your doctor can help find an infusion center. You can also call your local **Rx Onboarding and Care Coordinator (ROCC)** or Scholar Rock Supports at 1-833-777-5444, Monday – Friday, 8 AM – 8 PM ET.

To find a local ROCC, visit ScholarRockSupports.com.

Can I switch infusion locations?

Yes, you can switch infusion locations if your needs or preferences change, although some options may be restricted based on your insurance coverage. For help navigating the logistics or exploring other available sites, you can reach out to your ROCC. They can provide guidance and support to help ensure a smooth transition.

How will I know if ISEMBYLD is working?

If you're wondering if your treatment is working, you should ask your doctor or consult with your healthcare team. Be sure to discuss your ISEMBYLD treatment and share any changes you notice between infusions. Your doctor can help determine what those changes may indicate and whether any adjustments are needed.

What if I have more questions after my infusion?

Your ROCC and healthcare team are there to help. Don't hesitate to reach out if you have any additional questions. Ongoing communication with your healthcare team is an important part of your treatment experience.

> The Regional Clinical Specialist (RCS) team is also available to provide education and support around infusions whether performed in a hospital, home, or infusion center.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of ISEMBYLD?

- **Fractures.** Treatment with ISEMBYLD may increase the risk of bone fractures, including serious fractures. Bone fractures may happen with or without a fall or other injury. Your healthcare provider may consider stopping treatment with ISEMBYLD if you experience a bone fracture during treatment

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

Scholar Rock SUPPORTS™

Personalized help throughout your journey

Scholar Rock Supports was built around the idea that no one should have to navigate treatment alone.

Your **Rx Onboarding and Care Coordinator (ROCC)** can help with guiding you through insurance, financial assistance, infusion day logistics, caregiver support, and what to expect with ISEMBYLD.

Real Scholar Rock Supports dedicated team members

HAVE QUESTIONS ABOUT TREATMENT WITH ISEMBYLD?

Call Scholar Rock Supports at **1-833-777-5444,** Monday – Friday 8 AM – 8 PM ET.

FIND A LOCAL ROCC

To use the ROCC locator tool, visit **ScholarRockSupports.com**

ISEMBYLD™
(apitegromab-mstn) injection 150mg/3mL

Indication and Important Safety Information

What is ISEMBYLD (apitegromab-mstn)?

ISEMBYLD is a prescription medicine used to treat spinal muscular atrophy (SMA) in adults and children 2 years of age and older who are currently receiving a survival motor neuron 2 (SMN2)-targeted treatment.

IMPORTANT SAFETY INFORMATION

Before taking ISEMBYLD, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of low bone density or bone fractures
- are pregnant or plan to become pregnant. If you are pregnant or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. It is not known if ISEMBYLD will harm your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with ISEMBYLD
- are breastfeeding or plan to breastfeed. It is not known if ISEMBYLD passes into breast milk. Talk to your healthcare provider about the best way to feed your baby while on treatment with ISEMBYLD

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider, including your pharmacist, when you get a new medicine.

What are the possible side effects of ISEMBYLD?

- **Fractures.** Treatment with ISEMBYLD may increase the risk of bone fractures, including serious fractures. Bone fractures may happen with or without a fall or other injury. Your healthcare provider may consider stopping treatment with ISEMBYLD if you experience a bone fracture during treatment

The most common side effects of ISEMBYLD include:

- upper respiratory tract infections
- vomiting
- cough
- viral infections
- headache
- stomach flu (gastroenteritis)
- sore throat (pharyngitis)
- hypersensitivity

These are not all of the possible side effects of ISEMBYLD. Call your healthcare provider for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying full Prescribing Information and Patient Information.

ISEMBYLD™
(apitegromab-mstn) injection 150mg/3mL

IT'S TIME

For the first and only FDA-approved muscle-targeted treatment for SMA

We hope this brochure has helped you feel more comfortable with the ISEMBYLD infusion process. As you get ready for your first infusion, remember that Scholar Rock Supports and your **Rx Onboarding and Care Coordinators (ROCCs)** are always ready to support you throughout your treatment journey.

To learn more visit
ISEMBYLD.com

River
Treated with ISEMBYLD

IMPORTANT SAFETY INFORMATION

The most common side effects of ISEMBYLD include:

- upper respiratory tract infections
- vomiting
- cough
- viral infections
- headache
- stomach flu (gastroenteritis)
- sore throat (pharyngitis)
- hypersensitivity

These are not all of the possible side effects of ISEMBYLD. Call your healthcare provider for medical advice about side effects.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information.

ISEMBYLD and Scholar Rock Supports are trademarks of Scholar Rock.
Scholar Rock © 2026. All rights reserved. US-APIT-0353 09/26

Scholar Rock®